

CLAIMS


1. A composition comprising a compound comprising a formula $(M \equiv N)L^1$ and pharmaceutically acceptable salts thereof;

wherein N is nitrogen;

5 M is a transition metal; and

L^1 is a first crowned dithiocarbamate, wherein the first crowned dithiocarbamate comprises a first crown ether-containing group of formula $[(CH_2)_a-O]_b-(CH_2)_c$, wherein a is at least 2, b is at least 3, and c is at least 2.

10 2. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises a formula:



and pharmaceutically acceptable salt thereof;

wherein R^1 or R^2 comprises the first crown ether-containing group, or

15 R^1 and R^2 together comprise the first crown ether-containing group.

3. The composition of Claim 1, wherein the transition metal is a radioactive metal.


20 4. The composition of Claim 1, wherein the transition metal is ^{99m}Tc or ^{94m}Tc .

5. The composition of Claim 1, wherein the transition metal is ^{186}Re or ^{188}Re .

25


6. The composition of Claim 1, wherein subscript a in the formula of the first crown ether-containing group is 2, 3, 4, or 5.

7. The composition of Claim 1, wherein subscript a in the formula of the first crown ether-containing group is 2 or 3.
8. The composition of Claim 1, wherein subscript a in the formula of the first crown ether-containing group is 2.
9. The composition of Claim 1, wherein subscript b in the formula of the first crown ether-containing group is 3, 4, 5, 6, 7, or 8.
10. The composition of Claim 1, wherein subscript b in the formula of the first crown ether-containing group is 3, 4, 5 or 6.
11. The composition of Claim 1, wherein subscript c in the formula of the first crown ether-containing group is 2, 3, 4, or 5.
12. The composition of Claim 1, wherein subscript c in the formula of the first crown ether-containing group is 2 or 3.
13. The composition of Claim 1, wherein subscript c in the formula of the first crown ether-containing group is 2.
14. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises




15. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises

-71-




16. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises




5

17. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises




10

18. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises




- 15 19. The composition of Claim 1, wherein the first crowned dithiocarbamate comprises


-72-



20. The composition of Claim 1, wherein the first crowned dithiocarbamate is selected from the group consisting of




, and




5

21. The composition of Claim 1, wherein the compound is selected from the group consisting of:



10

-73-




- 5 22. The composition of Claim 1, wherein the compound further comprises L^2 and comprises a formula $(M \equiv N)L^1L^2$ and pharmaceutically acceptable salts thereof;

10 wherein L^2 is a second crowned dithiocarbamate, wherein the second crowned dithiocarbamate comprises a second crown ether-containing group of formula $[(CH_2)_a-O]_b-(CH_2)_c$, wherein a is at least 2, b is at least 3, and c is at least 2.

23. The composition of Claim 22, wherein the second crowned dithiocarbamate comprises a formula:

-74-



and pharmaceutically acceptable salt thereof;

wherein R¹ or R² comprise the second crown ether-containing group, or R¹ and R² together comprise the second crown ether-containing group.

5

24. The composition of Claim 22, wherein subscript a in the formula of the second crown ether-containing group is 2, 3, 4, or 5.

25. The composition of Claim 22, wherein subscript a in the formula of the second crown ether-containing group is 2 or 3.

10

26. The composition of Claim 22, wherein subscript a in the formula of the second crown ether-containing group is 2.

27. The composition of Claim 22, wherein subscript b in the formula of the second crown ether-containing group is 3, 4, 5, 6, 7, or 8.

15

28. The composition of Claim 22, wherein subscript b in the formula of the second crown ether-containing group is 3, 4, 5 or 6.

20

29. The composition of Claim 22, wherein subscript c in the formula of the second crown ether-containing group is 2, 3, 4, or 5.


30. The composition of Claim 22, wherein subscript c in the formula of the second crown ether-containing group is 2 or 3.

25


31. The composition of Claim 22, wherein subscript c in the formula of the second crown ether-containing group is 2.

-75-


32. The composition of Claim 22, wherein the second crowned dithiocarbamate comprises




5 33. The composition of Claim 22, wherein the second crowned dithiocarbamate comprises



34. The composition of Claim 22, wherein the second crowned
10 dithiocarbamate comprises




35. The composition of Claim 22, wherein the second crowned dithiocarbamate comprises




15

36. The composition of Claim 22, wherein the second crowned dithiocarbamate comprises

-76-




37. The composition of Claim 22, wherein the second crowned dithiocarbamate comprises




5

38. The composition of Claim 22, wherein the second crowned dithiocarbamate is selected from the group consisting of




, and



10


39. The composition of Claim 1, wherein the compound is selected from the group consisting of:

-77-




5

40. The composition of Claim 1, wherein the compound further comprises L^3 , L^4 , and L^5 and comprises a formula:



and pharmaceutically acceptable salts thereof;

10 wherein L^3 , L^4 , and L^5 each comprises an isonitrile of formula:



wherein q is 0 – 3;

Z is carbon or silicon;

R^3 , R^4 and R^5 are the same or different, and are selected from the group consisting of H, C_1 - C_{10} alkyl substituted with 0-5 R^6 , aryl

15


-78-

substituted with 0-5 R^6 , heteroaryl substituted with 0-5 R^6 , and macrocyclic crown ether containing 2-8 ether-oxygen atoms;

wherein R^6 is selected from the group consisting of H, OH, OR^7 , $C(=O)OR^7$, $C(=O)NR^8R^9$, $PO(OR^8)_2$, $PO(NR^8R^9)_2$ and SO_2R^7 ; and


R^7 , R^8 and R^9 are the same or different, and are selected from the group consisting of H, alkyl, aryl, and heteroaryl, or R^8 and R^9 together form a macrocyclic crown ether containing 2-8 ether-oxygen atoms.

41. The composition of Claim 1, wherein the compound further comprises L^3 , L^4 , and L^5 and comprises a formula:



and pharmaceutically acceptable salts thereof;

wherein L^3 , L^4 , and L^5 together form a tripodal chelator of formula:



wherein U is selected from the group consisting of $R^{13}B$, CR^{13} , and $P(=O)$;

A^1 , A^2 and A^3 are imine-N containing heterocycles;

A^4 , A^5 and A^6 are selected from the group consisting of NR^{14} , PR^{14} , S, and O;


R^{10} , R^{11} and R^{12} are selected from a group of formula $-(CH_2)_g-$, wherein g is 2-5;

R^{13} is selected from the group consisting of H, alkyl and aryl;

and


R^{14} is selected from the group consisting of H, alkyl, aryl, and alkoxyalkyl.

42. The composition of Claim 41, wherein the tripodal chelator is selected from the group consisting of



5

43. A composition comprising a compound comprising a formula:



and pharmaceutically acceptable salt thereof;


wherein M is a transition metal selected from the group consisting of Fe(II), Fe(III), Mn(II), Mn(III), Co(II), Co(III), Ni(II), Cu(II), Zn(II), Ru(II), Ru(III), Pd(II), and Pt(II);

p and p' are integers and are independently selected from 0-2;

10

R^1 and R^2 comprise a crown ether-containing group of formula $[(CH_2)_a-O]_b-(CH_2)_c$, wherein a is at least 2, b is at least 3, and c is at least 2, or wherein R^1 and R^2 together comprise the crown ether-containing group;

L^6 is a tripodal chelator with a formula selected from the group
5 consisting of



wherein U is selected from the group consisting of $R^{13}B$, CR^{13} , and $P(=O)$;

A^1 , A^2 and A^3 are imine-N containing heterocycles;

10 A^4 , A^5 and A^6 are selected from the group consisting of NR^{10} , PR^{10} , and S ;

R^{10} , R^{11} and R^{12} are selected from a group of formula $-(CH_2)_g-$, wherein g is 2-5;

R^{13} is selected from the group consisting of H , alkyl and aryl;

15 and

R^{14} is selected from the group consisting of H , alkyl, aryl, and alkoxyalkyl.

44. A method for radioimaging a subject comprising:

20

a) providing

i) a subject; and

ii) a composition comprising a compound comprising a formula $(M \equiv N)L^1$ and pharmaceutically acceptable salts thereof;

b) administering the composition to the subject; and

25

c) scanning at least a portion of the subject using a radioimaging device;

wherein N is nitrogen;

M is a radioactive transition metal; and

L^1 is a first crowned dithiocarbamate, wherein the first crowned dithiocarbamate comprises a first crown ether-containing group of formula $[(CH_2)_a-O]_b-(CH_2)_c$, wherein a is at least 2, b is at least 3, and c is at least 2.

5

45. The method of Claim 44, wherein at least a portion of the subject is tissue suspected of being diseased.

10

46. The method of Claim 44, wherein the at least a portion of the subject is myocardial tissue.

47. The method of Claim 44, wherein the subject is a mammal.

15


48. A method of treating a disease resulting from overproduction of nitric oxide or reactive oxygen species, comprising:

a) providing:

i) a subject with a disease; and

ii) a composition comprising a compound comprising a

formula:



20

and pharmaceutically acceptable salt thereof; and

b) administering the composition to the subject;

wherein M is a transition metal selected from the group consisting of Fe(II), Fe(III), Mn(II), Mn(III), Co(II), Co(III), Ni(II), Cu(II), Zn(II), Ru(II), Ru(III), Pd(II), and Pt(II);


25

p and p' are integers and are independently selected from 0-2;

R^1 and R^2 comprises a crown ether-containing group of formula $[(CH_2)_a-O]_b-(CH_2)_c$, wherein a is at least 2, b is at least 3, and c is at

least 2, or wherein R^1 and R^2 together comprise the crown ether-containing group;

L^6 is a tripodal chelator with a formula selected from the group consisting of:



wherein U is selected from the group consisting of $R^{13}B$, CR^{13} , and $P(=O)$;

A^1 , A^2 and A^3 are imine-N containing heterocycles;

A^4 , A^5 and A^6 are selected from the group consisting of NR^{10} , PR^{10} , and S;

R^{10} , R^{11} and R^{12} are selected from a group of formula $(CH_2)_g$, wherein g is 2-5;

R^{13} is selected from the group consisting of H, alkyl and aryl; and

R^{14} is selected from the group consisting of H, alkyl, aryl, and alkoxyalkyl.

49. A method of treating metal poisoning, comprising:

a) providing:

i) a subject with metal poisoning, and

ii) a composition comprising a crowned dithiocarbamate;

and

b) administering the composition to the subject;

wherein the crowned dithiocarbamate comprises a crown ether-containing group of formula $[(CH_2)_a-O]_b-(CH_2)_c$, wherein a is at least 2, b is at least 3, and c is at least 2.